



TITLE:

Studies on the Catalytic Action of AlCi

AUTHOR(S):

Tarama, Kimio; Kubota, Toshiaki

CITATION:

Tarama, Kimio ...[et al]. Studies on the Catalytic Action of AlCi. 京都大学
化学研究所報告 1950, 20: 59-59

ISSUE DATE:

1950-03-20

URL:


<http://hdl.handle.net/2433/74050>

RIGHT:

5. The apparent Arrhenius' activation energy of this reaction is 25 Kcal.

Kimio Tarama and Toshiaki Kubota.

From these results, the following mechanism is proposed, i.e. the reaction proceeds through the ionic intermediate complex between heptane and $\text{Al}_2\text{Cl}_{(6-n)}(\text{OH})_n$, produced from AlCl_3 and H_2O , in which n is smaller than 0.5 (simply represented by $\text{OHA}(\text{AlCl}_2)$).


$$\begin{array}{c} \text{CH}_3\text{-CH}_2\cdots\text{CH}_2\text{-}\overset{+}{\text{CH}}\text{-CH}_3 \\ | \\ \text{Cl} \\ | \\ (\text{Cl}\text{-Al}\text{-H})^- \\ | \\ \text{OH} \end{array}$$

form by the catalyst, mixture of CCl_4 and AlCl_3 , instead of H_2O .

(59)